

## List of Figures

Figure 1.1	A schematic research framework and thesis outline.	6
Figure 2.1	Correlation diagram between population density (logarithmic) and cumulative forest loss for Tropical Latin America, Tropical Africa and Tropical Asia. [After Pahari and Murai (1997)]	19
Figure 4.1	Schematic illustration of processes occurring after deforestation. [From Zhang <i>et al.</i> (1996b)]	47
Figure 4.2	Changes (i.e. difference between deforestation and control experiment) of (a) Hadley circulation [zonal average of vertical velocity from 180°W to 80°W ( $\text{Pa s}^{-1}$ )] and (b) Walker circulation [meridional average of vertical velocity from 9°S to 9°N ( $\text{Pa s}^{-1}$ )] in October. [From McGuffie <i>et al.</i> , (1995)]	57
Figure 5.1	Spatially averaged seasonal cycle of vertical velocity near the surface: (a) NCEP/NCAR Reanalysis (mean for 1976-96), and (b) control experiment, both for area bounded by 15°S to 30°N, 90°E to 130°E. (c) and (d) as for (a) and (b) respectively, but only for equatorial region, 15°S to 15°N, 90°E to 130°E.	67
Figure 5.2	Precipitation (mm) for January and July from observation (a and b) and simulation (c and d). Area with monthly precipitation greater than 200 mm is shaded. [Observations after Legates and Willmott (1990a)]	71
Figure 5.3	Surface air temperature (K) for January and July from observation (a and b) and simulation (c and d). [Observations after Legates and Willmott (1990b)]	72
Figure 5.4	(a) and (b): Observed (analysis) 1000 hPa wind vector ( $\text{ms}^{-1}$ ) with MSLP (hPa) for January and July (mean for 1976-96, after NCEP/NCAR Reanalysis), and (c) and (d): Simulated (control) 991 hPa wind vector ( $\text{ms}^{-1}$ ) for January and July, respectively.	74
Figure 5.5	Zonal (u) wind ( $\text{ms}^{-1}$ ) near surface: (a) and (b) are observed (analysis) fields at 1000 hPa level, mean for 1976-96 after NCEP/NCAR Reanalysis, and (c) and (d) are simulated (control) fields at 991 hPa level for January and July respectively.	75
Figure 5.6	As in Figure 5.5 but for meridional (v) wind ( $\text{ms}^{-1}$ ) near surface.	76

Figure 5.7	Seasonal variation of climatic changes over Southeast Asia after deforestation (areal average from 15°S to 20°N and from 95°E to 150°E): (a) total precipitation (mm month <sup>-1</sup> ), (b) total evapotranspiration (mm month <sup>-1</sup> ), (c) moisture divergence (mm month <sup>-1</sup> ), (d) surface net radiation (W m <sup>-2</sup> ), (e) sensible heat flux (Wm <sup>-2</sup> ), (f) surface air temperature (K). [From Zhang <i>et al.</i> (1996a)]	83
Figure 5.8	Changes of precipitation (mm month <sup>-1</sup> ) following deforestation for January and July (b).	85
Figure 5.9	Change of Walker circulation (meridional average of vertical velocity from 15°S to 15°N) (Pa/s) in January.	89
Figure 5.10	Change of Walker circulation (meridional average of vertical velocity from 15°S to 15°N) (Pa/s) in July.	90
Figure 5.11	Change of Hadley circulation (zonal average of vertical velocity from 90°E to 130°E) (Pa/s) in January.	91
Figure 5.12	Change of Hadley circulation (zonal average of vertical velocity from 90°E to 130°E) (Pa/s) in July.	92
Figure 5.13	Change of vertical velocity near surface (Pa/s) in January.	95
Figure 5.14	As in Figure 5.13 but for July.	96
Figure 5.15	Changes of divergence (s <sup>-1</sup> ) (shaded) and vorticity (s <sup>-1</sup> ) (white contour) near surface for January (associated streamlines are shown in the control and deforestation fields).	97
Figure 5.16	As in Figure 5.15 but for July.	98
Figure 6.1	An example of the vertical distribution of variables in a model with $N$ layers. The column of values on the right-hand side denotes the pressure at which variables are represented in a 15-level vertical resolution model, assuming a surface pressure of 1000hPa. [From Simmons and Bengtsson (1984)]	119
Figure 6.2	Standard 19-levels configuration for the Unified Model. The values labelled at some lower layers represent the $\sigma$ -level. [From Cullen (1993)]	119

Figure 6.3	Schematic illustration of the processes commonly included in atmospheric GCMs. The thickness of a particular arrow gives a qualitative indication of the importance of the interaction the arrow represents. [From Simmons and Bengtsson (1984)]	123
Figure 6.4	The Unified Model software system. [After Cullen (1993) and Met. Office (1996)]	133
Figure 8.1	Time series of global average precipitation (mm), surface evaporation (mm) and their difference for December - February, 91/92 and 92/93 of the control simulation.	174
Figure 8.2	Time series of global average precipitation (mm), surface evaporation (mm) and their difference for June - August, 1991 and 1994 of the control simulation.	175
Figure 8.3	Root depth (m) for the control (a) and deforestation (b) experiments. The inner box (b) is the study area which encloses the deforested region (i.e. grid-boxes with values).	179
Figure 8.4	As in Figure 8.3 but for snow-free albedo.	180
Figure 8.5	As in Figure 8.3 but for surface resistance (s/m).	181
Figure 8.6	As in Figure 8.3 but for roughness length (m).	182
Figure 8.7	As in Figure 8.3 but for surface capacity (mm).	183
Figure 8.8	As in Figure 8.3 but for vegetation fraction.	184
Figure 8.9	As in Figure 8.3 but for infiltration factor.	185
Figure 9.1	Precipitation (mm) for January and July from observation (a and b) and composite 10-case ensemble control simulations (c and d). [Observations after Legates and Willmott (1990a)]	195
Figure 9.2	Surface air temperature (K) for January and July from observation (a and b) and composite 10-case ensemble control simulations (c and d). [Observations after Legates and Willmott (1990b)]	195

Figure 9.3	(a) and (b): observed (analysis) 850 hPa wind vector ( $\text{ms}^{-1}$ ) with mean sea level pressure (hPa) for January and July [21-year mean (1976-96), after NCEP/NCAR Reanalysis], and (c) and (d): similar to (a) and (b), but for the control simulations. [composite from 10-case ensemble]	196
Figure 9.4	(a) and (b): observed (analysis) zonal (u) wind ( $\text{ms}^{-1}$ ) at 850 hPa level for January and July, respectively [21-year mean (1976-96)] after NCEP/NCAR Reanalysis. (c) and (d): simulated (control) fields at 850 hPa level for January and July respectively (composite from 10-case ensemble).	197
Figure 9.5	(a) and (b): observed (analysis) meridional (v) wind ( $\text{ms}^{-1}$ ) at 850 hPa level for January and July, respectively [21-year mean (1976-96)] after NCEP/NCAR Reanalysis. (c) and (d): simulated (control) fields at 850 hPa level for January and July respectively (composite from 10-case ensemble).	198
Figure 9.6	(a) and (b): observed 850 hPa velocity potential ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) for January and July, respectively [21-year mean (1976-96), after NCEP/NCAR Reanalysis]. (c) and (d): similar to (a) and (b), but for the control simulations (composite 10-case ensemble).	199
Figure 9.7	(a) and (b): observed 200 hPa velocity potential ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) for January and July, respectively [21-year mean (1976-96), after NCEP/NCAR Reanalysis]. (c) and (d): similar to (a) and (b), but for the control simulations (composite 10-case ensemble).	200
Figure 9.8	Standard deviation of monthly average vertical velocity from 10-case ensemble control simulations: (a) and (b) are meridional average (from $15^\circ\text{S}$ to $15^\circ\text{N}$ ) in January and July respectively; and (c) and (d) are zonal average ( $90^\circ\text{E}$ to $150^\circ\text{E}$ ) in January and July respectively.	207
Figure 9.9	Standard deviation of monthly precipitation from the 10-case ensemble: (a) and (b) for the control in January and July, respectively; (c) and (d) for the deforestation in January and July, respectively.	208
Figure 9.10	Changes in surface temperature (K) in January (a) and July (b), respectively, with local changes significant at the 5% level shaded; (c) and (d): associated grid-point values of surface temperature changes in the study area for January and July, respectively.	219
Figure 9.12	Change in surface evaporation (mm) in January (a) and July (b). Areas in which changes are locally significant at the 5% level are shaded.	221

Figure 9.13	Change in 1000 hPa relative humidity (%) in January (a) and July (b). Areas in which changes are locally significant at the 5% level are shaded.	222
Figure 9.14	Change in net down surface short-wave radiation ( $\text{Wm}^{-2}$ ) in January (a) and July (b). Areas in which changes are locally significant at the 5% level are shaded.	223
Figure 9.15	Change in net down surface long-wave radiation ( $\text{Wm}^{-2}$ ) in January (a) and July (b). Areas in which changes are locally significant at the 5% level are shaded.	224
Figure 9.16	Change in net down surface radiation ( $\text{Wm}^{-2}$ ) in January (a) and July (b). Areas in which changes are locally significant at the 5% level are shaded.	225
Figure 9.17	Conceptual model of the atmospheric moisture fluxes over a land region. Terms $P_m$ and $P_a$ are precipitation of local evaporative and advective origin, respectively. Modified from Brubaker <i>et al.</i> (1993), with $R_s$ and $R_d$ are added.	237
Figure 10.1	Change of Walker circulation (meridional average of vertical velocity from $15^\circ\text{S}$ to $15^\circ\text{N}$ ) ( $\text{Pa/s}$ ) in January (composite from 10-case ensemble).	245
Figure 10.2	Change of Walker circulation (meridional average of vertical velocity from $15^\circ\text{S}$ to $15^\circ\text{N}$ ) ( $\text{Pa/s}$ ) in July (composite from 10-case ensemble).	246
Figure 10.3	Change of Hadley circulation (zonal average of vertical velocity from $90^\circ\text{E}$ to $150^\circ\text{E}$ ) ( $\text{Pa/s}$ ) in January (composite from 10-case ensemble).	247
Figure 10.4	Change of Hadley circulation (zonal average of vertical velocity from $90^\circ\text{E}$ to $150^\circ\text{E}$ ) ( $\text{Pa/s}$ ) in July (composite from 10-case ensemble).	248
Figure 10.5	Changes in mean sea level pressure (hPa) in January and July. No changes are locally significant at the 5% level.	253
Figure 10.6	Change in vertical velocity pattern at lower level (850 hPa) ( $\text{Pa/s}$ ) in January. Changes locally significant at the 5% level are shaded.	254

Figure 10.7	Change in vertical velocity pattern at lower level (850 hPa) (Pa/s) in July. Changes locally significant at the 5% level are shaded.	255
Figure 10.8	Change in velocity potential at lower level (850 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in January. Arrows indicate divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	258
Figure 10.9	Change in velocity potential at upper level (200 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in January. Arrows indicate divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	259
Figure 10.10	Change in velocity potential at lower level (850 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in July. Arrows indicate divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	260
Figure 10.11	Change in velocity potential at upper level (200 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in July. Arrows indicate divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	261
Figure 10.12	Change in stream function at middle level (500 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in January. Arrows indicate non-divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	263
Figure 10.13	Change in stream function at middle level (500 hPa) ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in July. Arrows indicate non-divergent wind vector ( $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded.	264
Figure 10.14	The meridional distribution of surface moisture flux, precipitation and moisture convergence (mm) in January for the control (solid line) and deforestation (dotted line) (zonally averaged from $90^\circ\text{E}$ to $150^\circ\text{E}$ , 10-case ensemble).	275
Figure 10.15	The meridional distribution of surface moisture flux, precipitation and moisture convergence (mm) in July for the control (solid line) and deforestation (dotted line) (zonally averaged from $90^\circ\text{E}$ to $150^\circ\text{E}$ , 10-case ensemble).	276
Figure 10.16	The zonal distribution of surface moisture flux, precipitation and moisture convergence (mm) in January for the control (solid line) and deforestation (dotted line) (meridionally averaged from $10^\circ\text{S}$ to $20^\circ\text{N}$ , 10-case ensemble).	277

Figure 10.17	The zonal distribution of surface moisture flux, precipitation and moisture convergence (mm) in July for the control (solid line) and deforestation (dotted line) (meridionally averaged from 10°S to 20°N, 10-case ensemble).	278
Figure 10.18	The meridional distribution of net down surface radiation ( $\text{Wm}^{-2}$ ) in January for the control (solid line) and deforestation (dotted line) (zonally averaged from 90°E to 150°E, 10-case ensemble).	279
Figure 10.19	The meridional distribution of net down surface radiation ( $\text{Wm}^{-2}$ ) in July for the control (solid line) and deforestation (dotted line) (zonally averaged from 90°E to 150°E, 10-case ensemble).	280
Figure 10.20	Meridional distribution (zonally averaged from 90°E to 150°E) of boundary layer entropy in terms of boundary layer potential temperature, $\Pi$ , in Kelvins for the control (solid line) and deforestation (dotted line): (a) for January, and (b) for July. (composite from 10-case ensemble).	281
Figure 10.21	Change in boundary layer potential temperature, $\Pi$ , (K) in January (a) and July (b). Changes locally significant at the 5% level are shaded (composite from 10-case ensemble).	282
Figure 10.22	A mechanism relating Southeast Asian deforestation and the strengthening of the large scale monsoon circulation.	286
Figure 10.23	Velocity potential ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in January at 850 hPa (a) and 200 hPa (b), and changes in January at 850 hPa (c) and 200 hPa (d). Arrows represent its associated divergent flow (i.e. divergent wind vector in $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded (composite from 10-case ensemble).	290
Figure 10.24	Velocity potential ( $\times 10^6 \text{ m}^2\text{s}^{-1}$ ) in July at 850 hPa (a) and 200 hPa (b), and changes in January at 850 hPa (c) and 200 hPa (d). Arrows represent its associated divergent flow (i.e. divergent wind vector in $\text{ms}^{-1}$ ). Changes locally significant at the 5% level are shaded (composite from 10-case ensemble).	291