# LIST OF TABLES

Page

## **CHAPTER 3:**

Table 3.1. Spatially incomplete network of daily data stations for precipitation. The period of records for all thestationsisfrom1931to2001.\*metersabovesealevel.32

Table 3.2. Spatially incomplete network of daily data stations for precipitation. The period of records for all thestationsisfrom1931to2001.\*metersabovesealevel.37

# **CHAPTER 4:**

Table 4.1. Total annual precipitation PCA resulting regions identified according to the known Mexicanclimatology(García,80

Table 4.2. Wet season precipitation PCA resulting regions identified according to the known Mexicanclimatology(García,87

Table 4.3. Dry season precipitation PCA resulting regions identified according to the known Mexicanclimatology(García,94

Table 4.4. List of stations with monthly mean temperature. The period of records for all the stations is from 1941to2001.\*metersabovesealevel.96

XV

#### **CHAPTER 5:**

Table 5.1. List of temperature stations (with data from 1941 to 2001) for which weather extreme indices showcorrelations (Kendall's tau) that are statistically significant at the 5 and 1% level. \* Stations numbers are incorrespondencewithTable3.2andFig.3.6.113

Table 5.2. List of temperature stations (with data from 1941 to 2001) for which weather extreme indices show correlations (Kendall's tau) between the precipitation extreme indices with time that are no statistically significant. \* Stations numbers are in correspondence with Table 3.2 and Fig. 3.6. 113

 Table. 5.3. Geographical patterns for rainfall extreme indices following positive/negative correlations using

 Kendall's tau and statistical significant levels at 5 and 1%. The number of cases are classified defining the Tropic

 of
 Cancer
 as
 the
 divide
 to
 separate
 the
 northern/southern
 regions.

 121

Table. 5.4. Geographical patterns following wet/dry conditions. The numbers of cases are classified defining theTropicofCancerasthedividetoseparatenorthern/southernresults.122

## **CHAPTER 6:**

Table 6.1. List of temperature stations (with data from 1941 to 2001) for which weather extreme indices show correlations (Kendall's tau) that are statistically significant at 5 and 1% level. The stations with the most statistically significant correlations are marked with (11), (8), and (7) depending on the number. \* Stations numbers are in correspondence with Table 3.2 and Fig. 3.6. 136

Table 6.2. List of temperature stations (with data from 1941 to 2001) show correlations (Kendall's tau) betweenthe precipitation extreme indices with time, that are not statistically significant. \* Stations numbers are incorrespondencewithTable3.2andFig.137

Table 6.3. Geographical patterns of temperature following positive/negative correlations of the extreme temperature indices using Kendall's tau and statistical significant levels at 5 and 1%. The number of cases is classified defining the Tropic of Cancer as the limit to separate the northern/southern regions. 152

 Table 6.4. Geographical patterns of temperature following warm/cold conditions of the extreme temperature indices using kendall's tau and statistical significant levels at 5 and 1%. The number of cases is classified defining the Tropic of Cancer as the limit to separate the northem/southern regions.

 155

## **CHAPTER 7:**

Table 7.1. Lag cross-correlations between the standardised versions of Regional Precipitation Averages and thedifferent ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El NiñoIndex (MEI). The linear correlation is calculated using the Pearson function. Lags (leads) are expressed inmonths and related to the maximum correlation found after trying several lags and leads. Regions displayed herearedefinedinTable4.1.196

Table 7.2. Lag cross-correlations between the RX1day (Max 1-day Precipitation) Index and the different

standardised versions of the ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the maximum correlation found after trying several lags and leads. 199

 Table 7.3. Lag cross-correlations between the RX5day (Max 5-day Precipitation) and the different ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the maximum correlation found after trying several lags and leads.

 201

 Table 7.4.
 Lag cross-correlations between the DTR (Daily Temperature Range) and the different ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the maximum correlation found after trying several lags and leads.

 202

 Table 7.5. Lag cross-correlations between the TN10P (Cool Night Frequency) and the different ENSO indices:

 the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear

 correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the

 maximum
 correlation

 found
 after

 trying
 several

 lags
 and

 leads.

 204

Table 7.6. Lag cross-correlations between the TN90P (Hot Night Frequency) and the different ENSO indices:the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linearcorrelation is calculated using the Pearson function. Lags (leads) are expressed in months and related to themaximumcorrelationfoundaftertryingseverallagsandleads.206

Table 7.7. Lag cross-correlations between the TNn (Coolest night) and the different ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation is

calculated using the Pearson function. Lags (leads) are expressed in months and related to the maximum correlation found after trying several lags and leads. 208

Table 7.8. Lag cross-correlations between the TNx (Hottest night) and the different ENSO indices: the SouthernOscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation iscalculated using the Pearson function. Lags (leads) are expressed in months and related to the maximumcorrelationfoundaftertryingseverallagsandleads.210

 Table 7.9. Lag cross-correlations between the TX10P (Cool day frequency) and the different ENSO indices: the

 Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear

 correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the

 maximum
 correlation

 found
 after

 trying
 several

 lags
 and

 leads.

 212

 Table 7.10. Lag cross-correlations between the TX90P (Hot day frequency) and the different ENSO indices: the

 Southern
 Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear

 correlation is calculated using the Pearson function. Lags (leads) are expressed in months and related to the

 maximum
 correlation

 found
 after

 trying
 several

 lags
 and

 leads.
 214

Table 7.11. Lag cross-correlations between the TXn (Coolest day) and the different ENSO indices: the SouthernOscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation iscalculated using the Pearson function. Lags (leads) are expressed in months and related to the maximumcorrelationfoundaftertryingseverallagsandleads.216

Table 7.12. Lag cross-correlations between the TXx (Hottest day) and the different ENSO indices: the Southern Oscillation Index (SOI), El Niño 3.4 index, and the Multivariate El Niño Index (MEI). The linear correlation is

calculated using the Pearson function. Lags (leads) are expressed in months and related to the maximum correlation found after trying several lags and leads. 218